arXiv:1402.5268v1 [math.NT] 21 Feb 2014

SOLUTION OF THE DIOPHANTINE EQUATION $x_1x_2x_3\cdots x_{m-1}=z^n$

ZAHID RAZA, HAFSA MASOOD MALIK

Abstract. This work determine the entire family of positive integer solutions of the diophantine equation. The solution is described in terms of $\frac{(m-1)(m+n-2)}{2}$ or $\frac{(m-1)(m+n-1)}{2}$ positive parameters depending on n even or odd. We find the solution of a diophantine system of equations by using the solution of the diophantine equation. We generalized all the results of the paper [5].

1. Introduction and Preliminaries

All the solutions of a diophantine equation of the form

$$ax-by=c,$$

has been found. But the theory on this equation in the literature can not apply on a diophantine equation of the form

$$x_1x_2x_3\cdots x_{m-1}=z^n. \tag{1.1}$$

So, we archived the solution of the equation 1.1. In [5], the author worked on the diophantine equation1.1 for $m=3$ with $n=2,3,4,5,6$, and $m=4$ with $n=2$; furthermore, he also worked on a diophantine system of $2-$equation of $5-$variables. We extent all these results to the general case that is for all $m\geq 3$ and $n\geq 2$ and use it to find the solution of a system of s-diophantine equation in t variables.

The authors have not been able to find material on the equations of this paper. However, W. Sierpinski's book, Elementary Theory of Numbers listed two papers with material on equations of the form, $x_1x_2x_3\cdots x_n=t^k$. One is a 3-page paper published in 1955 (see reference[3]). And the other, a 10-page paper published in 1933 (see reference[4]). The authors have not been able to access these two papers. Even if some of the results in those two works; overlap with some of the results in this work, it is quite likely, that the methods used in this paper; will be different from those used in the above mentioned papers.

Key words and phrases. Divisor; diophantine equation; diophantine system of equations; the greatest common divisor.

- An integer b is called divisible by an other integer $a \neq 0$, if there exist some integer c such that
$$b = ac.$$
Symbolically,
$$a \mid b.$$
- Let a and b be given integer, with at least one of them different from zero. A positive integer d is called the greatest common divisor of the integers a, b ???
$\diamondsuit$ If $d \mid a$ and $d \mid b$.
$\diamondsuit$ Whenever there is c such that $c \mid a$ and $c \mid b$, then $c \leq d$.
Symbolically, $d = \gcd(a, b)$
- Two integer a, b are said to be relatively prime if $\gcd(a, b) = 1$.
- Whenever $\gcd(a, b) = d$; then $\gcd(\frac{a}{d}, \frac{b}{d}) = 1$.

The first theorem, well-known and widely used in number theory, is known as Euclids and for a proof see reference [1] or reference [2].

Theorem 1.1. *Assume that α, β, γ are positive integer such that $\gcd(\alpha, \beta) = 1$, and $\alpha\beta \mid \gamma$, then $\alpha \mid \gamma$.*

The following theorem can be proved by using the fundamental theorem of arithmetic. It can also be proved without the use of the fundamental theorem (for example [1]).

Theorem 1.2. *Consider that α, β and n to be positive integers. If $\alpha^n \mid \beta^n$, then $\alpha \mid \beta$*

In reference [1], the reader can find a proof of theorem 1.3 that makes use of theorem 1.1 and 1.2 ; but not the factorization theorem of a positive integer into prime powers.

Theorem 1.3. *If $\eta, \alpha, \beta, \gamma$ and n are positive integer such that*
$$\alpha\beta = \eta\gamma^n \quad \textit{where} \quad \gcd(\alpha, \beta) = 1,$$
then
$$\alpha = \sigma{\alpha'_1}^n, \quad \beta = \varsigma{\beta'_1}^n, \quad \eta = \varsigma\sigma \quad \textit{and} \quad \gamma = \alpha'_1\beta'_1$$
where, $\gcd(\alpha'_1, \beta'_1) = 1 = \gcd(\sigma, \varsigma)$.

Theorem 1.4. *Let k be a positive integer, and assume the 3-variables diophantine equation*
$$xy = kz^n.$$
Then all positive integer solutions can be described by 2-parametric formulas;
$x = k_1{t_1}^n, \quad y = k_2{t_2}^n, \quad z = t_1t_2$ *where,* $\gcd(k_1, k_2) = 1 = \gcd(t_1, t_2)$.

2. Main Result

Theorem 2.1. *The diophantine equation in m-variables*

$$x_1x_2x_3\cdots x_{m-1} = z^n$$

is equivalent to the $m+3$-variable system of equations

$$X_{m-1}X_{m-2}x_1x_2\cdots x_{m-3} = v{Z_0}^n, \tag{2.1}$$

$$w^{n-2} = vd^2 \tag{2.2}$$

where $d, X_{m-1}X_{m-2}, Z_0, w, v$ are positive integer variable such that
$x_{m-1} = \theta X_{m-1}, \quad x_{m-2} = \theta X_{m-2}, \quad z = wZ_0, \quad \theta = wd,$
$\theta = \gcd(x_{m-1}, x_{m-2}), \quad w = \gcd(z, \theta), \gcd(X_{m-1}, X_{m-2}) = 1 = \gcd(Z_0, d).$

Proof. Consider the equation 1.1 and let

$$\theta = \gcd(x_{m-1}, x_{m-2})$$

such that

$$\Big\{x_{m-1} = \theta X_{m-1}; x_{m-2} = \theta X_{m-2}, \gcd(X_{m-1}, X_{m-2}) = 1\Big\}, \tag{2.3}$$

then from equation 1.1 and 2.3 we obtain,

$$\theta^2 X_{m-1}X_{m-2}x_1x_2\cdots x_{m-3} = z^n. \tag{2.4}$$

Assume that $w = \gcd(z, \theta)$, then

$$\Big\{z = wZ_0;\ \theta = wd,\ \gcd(Z_0, d) = 1\Big\} \tag{2.5}$$

and from equation 2.11 and 2.4 we further have,

$$w^2d^2X_{m-1}X_{m-2}x_1x_2\cdots x_{m-1} = Z_0^n w^n, d^2X_{m-1}X_{m-2}x_1\cdots x_{m-3} = Z_0^n w^{n-2}.$$

Thus

$$d^2X_{m-1}X_{m-2}x_3\cdots x_{m-3} = Z_0^n w^{n-2}, \quad n \geq 2 \tag{2.6}$$

since $\gcd(d, Z_0) = 1$, it follows that $\gcd(d^2, Z_0^n) = 1$ which together with equation 2.4 and theorem 1.1; implies that d^2 must be a divisor of w^{n-2} that is

$$w^{n-2} = vd^2, \quad \text{for some} \quad v \in \mathbb{Z} \tag{2.7}$$

from equations 2.7 and 2.6, we have the result. □

Theorem 2.2. *All the positive solution of the equation 2.2 can be describe by the parameters as:*

i: *If n is odd, then*

$$w = \prod_{i=0}^{\frac{n-3}{2}} r_{2i+1}g^2, \quad d = \prod_{i=0}^{\frac{n-3}{2}} (r_{2i+1})^i g^{n-2}, \quad v = \prod_{i=0}^{\frac{n-3}{2}} (r_{2i+1})^{n-2i-2}.$$

ii: *If n is even, then*

$$w=\prod_{i=0}^{\frac{n-4}{2}} r_{2i+1}h,\ \ d=\prod_{i=0}^{\frac{n-4}{2}}(r_{2i+1})^i h^{\frac{n-2}{2}},\quad v=\prod_{i=0}^{\frac{n-4}{2}}(r_{2i+1})^{n-2i-2}.$$

Proof. (i) : If n is odd and let $D_0=\gcd(w,d)$, then

$$w=D_0.r_1 \text{ and } d=D_0.r_0 \text{ such that } \gcd(r_1,r_0)=1. \tag{2.8}$$

So from 2.1 and 2.2, we have $D_0^{n-2}r_1^{n-2}=vD^2r_0^2$

$$D_0^{n-4}{r_1}^{n-2}=vr_0^2 \tag{2.9}$$

Since $\gcd(r_1,r_0)=1$, so using theorem 1.1, we have $r_0^2|D_0^{n-4}$. Consider $D_1=\gcd(D_0,r_0)$, then

$$D_0=D_1r_3 \text{ and } r_0=D_1r_2 \text{ such that } \gcd(r_3,r_2)=1. \tag{2.10}$$

Thus ${D_1}^{n-4}{r_3}^{n-4}r_1^{n-2}=v{D_1}^2{r_2}^2$

$${D_1}^{n-6}{r_3}^{n-4}r_1^{n-2}=v{r_2}^2 \tag{2.11}$$

where $\gcd(r_2,r_1r_3)=1$ using theorem1.1, we get ${r_2}^2 \mid {D_1}^{n-6}$. Continued in this way and let $D_i=\gcd(r_{2i-2},D_{i-1})$, then

$$D_{i-1}=D_ir_{2i+1} \text{ and } r_{2i-2}=D_ir_{2i} \text{ such that } \gcd(r_{2i+1},r_{2i})=1 \tag{2.12}$$

we obtain $D_i^{n-2i}{r_{2i+1}}^{n-2i}{r_{2i-1}}^{n-2i-6}\cdots{r_3}^{n-2}r_1^{n-2}=vD_i^2.{r_{2i}}^2$

$$D_i{}^{n-2i-2}{r_{2i+1}}^{n-2i}{r_{2i-1}}^{n-4i-6}\cdots{r_3}^{n-2}r_1^{n-2}=v{r_{2i}}^2. \tag{2.13}$$

Since $\gcd(r_{2i},r_1r_3\ldots r_{2i+1})=1$ and using theorem 1.1, we have ${r_{2i}}^2 \mid {D_i}^{n-2i-4}$ where $i=0,1,2,\ldots,\frac{n-5}{2}$. Finally we have ${r_{n-5}}^2 \mid D_{\frac{n-5}{2}}$ as n is odd then $D_{\frac{n-5}{2}}={r_{n-2}{r_{n-5}}^2}$, $v=r_{n-2}r_{n-4}^3r_{n-6}^5\cdots r_3^{n-4}r_1^{n-2}$ $D_{\frac{n-5}{2}}=r_{n-2}{r_{n-5}}^2$ and $r_{n-7}={r_{n-2}{r_{n-5}}^3}$
substituting backward we get the required result, where $g=r_{n-5}$ and $h=r_{n-6}$. □

Remark **2.3.**

- When n is an odd integer $\theta=\prod_{i=0}^{\frac{n-3}{2}}(r_{2i+1})^{i+1}g^n$
- When n is an even integer $\theta=\prod_{i=0}^{\frac{n-4}{2}}(r_{2i+1})^{i+1}h^{\frac{n}{2}}$

Theorem 2.4. *Let us consider the m-variables Diophantine equation 1.1.*

i: *If n is odd, then all the positive solution of this equation can be describe by the parametric formulas*

$$x_j=\prod_{i=0}^{\frac{n-3}{2}}\left(k_{2i+1}^{j-1}\right)^{n-2i-2}\left(\prod_{t=1}^{j-1}\left(\gamma_t^{j-t}\right)\prod_{t=1}^{j}\left(\gamma_j^{j-t}\right)\right)^{n-1}\gamma_j^{j-t}\eta_j^{m-2-j}$$

$j=1,2,\ldots,m-3$

$$x_{m-2} = (\prod_{t=1}^{m-3}(\prod_{i=0}^{\frac{n-3}{2}}((k_{2i+1}^{t-1}l_{2i+1}^{m-3})^{i+1})(k_{2i+1}^{m-3})^{n-2i-2})(\gamma_t^{m-2-t})^{n-1})s_2^n g^n$$

$$x_{m-1} = (\prod_{t=1}^{m-3}(\prod_{i=0}^{\frac{n-3}{2}}((k_{2i+1}^{t-1}l_{2i+1}^{m-3})^{i+1})l_{2i+1}^{m-3^{n-2i-2}})(\eta_t^{m-2-t})^{n-1})s_1^n g^n$$

$$z = \prod_{\lambda=1}^{m-3}\prod_{i=0}^{\frac{n-3}{2}}(k_{2i+1}^{\lambda-1}l_{2i+1})(\prod_{i=1}^{m-3}\prod_{\lambda=0}^{m-2-i}\eta_i\gamma_i^{\lambda})s_1s_2g^2$$

ii: *If n is even, then all the positive solution of this equation can be describe by the parametric formulas*

$$x_j = \prod_{i=0}^{\frac{n-4}{2}}\left(k_{2i+1}^{j-1}\right)^{n-2i-2}\left(\prod_{t=1}^{j-1}\left(\gamma_t^{j-t}\right)\prod_{t=1}^{j}\left(\gamma_j^{j-t}\right)\right)^{n-1}\gamma_j^{j-t}\eta_j^{m-2-j}$$

for all $j = 1, 2, \ldots, m-3$

$$x_{m-2} = (\prod_{t=1}^{m-3}(\prod_{i=0}^{\frac{n-3}{2}}((k_{2i+1}^{t-1}l_{2i+1}^{m-3})^{i+1})(k_{2i+1}^{m-3})^{n-2i-2})(\gamma_t^{m-2-t})^{n-1})s_2^n h^{\frac{n}{2}}$$

$$x_{m-1} = (\prod_{t=1}^{m-3}(\prod_{i=0}^{\frac{n-4}{2}}((k_{2i+1}^{t-1}l_{2i+1}^{m-3})^{i+1})l_{2i+1}^{m-3^{n-2i-2}})(\eta_t^{m-2-t})^{n-1})s_1^n h^{\frac{n}{2}}$$

$$z = \prod_{\lambda=1}^{m-3}\prod_{i=0}^{\frac{n-3}{2}}(k_{2i+1}^{\lambda-1}l_{2i+1}^{m-3})(\prod_{i=1}^{m-3}\prod_{\lambda=0}^{m-2-i}\eta_i^{\lambda}\gamma_i^{\lambda})s_1s_2g^2$$

where $k_{2i+1}^t, l_{2i+1}^{m-3}, b_i^t a_i^t, s_1, s_2, h$ *and* g *are positive integers such that* $1 = \gcd(s_1, s_2)$, $\gcd(k_{2i-1}, k_{2i-1}^t l_{2i-1}^{m-3}) = 1 = \gcd(\gamma_i^t\eta_i, \gamma_i) = 1 \ \forall\ i = 1, 2, \ldots, \frac{n-6}{2},\ t = 1, 2, \ldots, m-3.$

Proof. Since $n \geq 2$, then by theorem 2.1, the given equation is equivalent to the diophantine system, $v = a_0$, $X_{m-1}X_{m-2}x_1x_2x_3....x_{m-3} = Z_0^n a_0$ and $w^{n-2} = vd^2$. Thus by theorem 2.2 and remark 2.3 let $P_1 = \gcd(x_1, Z_0)$, then $x_1 = P_1.X_1$ and $Z_0 = P_1.Z_1$, so $X_{m-1}X_{m-2}X_1x_2\cdots x_{m-3} = Z_1^n P_1^{n-1}a_0$.
Again let $\gcd(Z_1, X_1) = 1$, then, we have $X_1 \mid a_0P_1^{n-1} \Rightarrow X_1.a_1 = a_0P_1^{n-1}$ and by theorem 1.3 $X_1 = \alpha_1\gamma_1^{n-1}$ and $a_1 = \beta_1\eta_1^{n-1}$ such that $\alpha_1\beta_1 = a_0$ and $\gamma_1\eta_1 = P_1$ where $\gcd(\alpha_1, \beta_1) = 1 = \gcd(\gamma_1, \eta_1)$.
We have $X_{m-1}X_{m-2}x_2x_3....x_{m-3} = Z_0^n a_1$, continued in this way and let $P_i = \gcd(x_i, Z_{i-1})$. then $x_i = P_i.X_i$ and $Z_{i-1} = P_i.Z_i$. Thus $X_{m-1}X_{m-2}X_i\prod_{j=i+1}^{m-3}x_j = Z_i^n P_i^{n-1}a_{i-1}$. Since $\gcd(Z_i, X_i) = 1$ then, $X_i \mid a_{i-1}P_i^{n-1} \Rightarrow X_i.a_i = a_{i-1}P_i^{n-1}$ so by theorem 1.3, we get $X_i = \alpha_i\gamma_i^{n-1}$ and $a_i = \beta_i\eta_i^{n-1}$ such that $\alpha_i\beta_i = a_i$ and $\gamma_i\eta_i = P_i$ where $\gcd(\alpha_i, \beta_i) = 1 = \gcd(\gamma_i, \eta_i)$. We have $X_{m-1}X_{m-2}\prod_{j=i+1}^{m-3}x_j = Z_{i-1}^n a_i \quad \forall\ i = 1, 2, 3, \ldots, m-3$ and at last we get $X_{m-1}X_{m-2} = Z_{m-3}^n a_{m-3}$ then by theorem 1.3 $X_{m-2} = \alpha_{m-2}s_1^n$ and $X_{m-1} = \beta_{m-2}s_2^n$ such that $\alpha_{m-2}\beta_{m-2} = a_{m-3}$ and $s_1s_2 = Z_{m-4}$

with $\gcd(\alpha_{m-2}, \beta_{m-2}) = 1 = \gcd(s_1, s_1)$. Now by using theorem 1.3 and technique in proof of theorem 2.2, we have for all $j = 1, 2, 3, \ldots, m-2$
$\alpha_j = (\prod_{i=1}^{m-3} (k_{2i+1}^{j-1})^{n-2i-2})(\prod_{t=1}^{j} \gamma_t^{j-t})^{n-1}, \quad \beta_j = (\prod_{i=1}^{m-3} (l_{2i+1}^{j-1})^{n-2i-2})(\prod_{t=1}^{j} \eta_t^{j-t})^{n-1}.$
Put in equation2.2 and 2.4 we get required result. □

Remark **2.5.** The solution is described in terms of $\frac{(m-1)(m+n-2)}{2}$ or $\frac{(m-1)(m+n-1)}{2}$ positive parameters depending on n even or odd.

Let $p_1, p_2, \ldots, p_s$ and r be positive integers and suppose that $t = p_1 + p_2 + \cdots + p_s - r$

Theorem 2.6. *Consider the t-variables diophantine system of s equations,*
$x_{11}x_{12}x_{13}....x_{1p_1-1} = z_1^{k_1}$
$x_{21}x_{22}x_{23}....x_{2p_2-1} = z_2^{k_2}$
$\vdots$
$x_{i1}x_{i2}x_{i3}....x_{ip_i-1} = z_i^{k_i}$
$\vdots$
$x_{s1}x_{s2}x_{s3}....x_{sp_s-1} = z_s^{k_s}$ *where* $2 \leq k_j'$ *s are positive number where s no.of equations and r is no of repeated variables in this system. Then all the positive solutions of this system of equations can be describe by using the following algorithm.*

Proof.
- Write solution of each equation by using theorem 2.4
- Select one variable from r-repeated variables and find the unique d in solution by using technique theorem 2.4
- Replace those values of parameters appear in selected repeated, in other variable.
- Do the same activity with other repeated variable.
- If all repeated variables have unique solution then substituting the values of parameter that exist in that variables in other variable.

□

To explain above theorem, here is two illustration below

Example 2.7. *Consider the 6-variables Diophantine system,* $x_1x_2x_3 = z_1^3$ *and* $x_3x_4 = z_2^2$. *Then all the positive solutions of this system equations can be describe by* 11 *parametric formulas*
$x_1 = ak_1'^2 l_1' \gamma_1'^2 R_1^3 g^3, \quad x_2 = ak_1' l_1'^2 \eta_1'^2 R_2^3 g^3, \quad x_3 = a\gamma_1' \eta_1' b_1^3 f^6, \quad x_4 = a\gamma_1' \eta_1' b_1^3 r_1^2,$
$z_1 = al_1' k_1' \gamma_1' \eta_1' b_1 f^2 R_1 R_2 g^2, \quad z_2 = a\gamma_1' \eta_1' b_1^3 f^3 r_1.$
Solution:-
Step 1: We apply theorem 2.4 to get the following solution for equation 1 *of the system*
$\{x_1 = k_1 k_1'^2 l_1' \gamma_1'^2 R_1^3 g^3, \ x_2 = k_1 k_1' l_1'^2 \eta_1'^2 R_2^3 g^3, \ x_3 = k_1 \gamma_1^3 \gamma_1' \eta_1', \ z_1 = l_1' k_1' k_1 \gamma_1 \gamma_1' \eta_1' R_1 R_2 g^2\}$
and for equation 2 *of the system*

$\left\{x_4 = dr_1^2, \quad x_3 = r_2^2 d, \quad z_2 = dr_1 r_2\right\}$ *Step 2: Since* x_3 *is the repeated variable in the equations of the system, so*

$$r_2^2 d = x_3 = k_1 \gamma_1^3 \gamma_1' \eta_1' \tag{2.14}$$

Now we will find that unique d *for* x_3*, and suppose* $a = \gcd(d, k_1)$ *such that*

$$\left\{d = aD_1; k_1 = aK_1, gcd(D_1, K_1) = 1\right\}$$

put in equation 2.14

$$r_2^2 D_1 = \gamma_1^3 \gamma_1' \eta_1' K_1 \ \Rightarrow D_1 \mid \gamma_1^3 \gamma_1' \eta_1' \tag{2.15}$$

then there exist a positive integer b *such that* $D_1 b = \gamma_1^3 \gamma_1' \eta_1'$ *by using theorem 2.4* $D_1 = \beta_1 c_1^3, \quad b = \beta_2 c_2^3, \quad \gamma_1 = c_1 c_2, \quad \gamma_1' \eta_1' = \beta_1 \beta_2$ *such that* $\gcd(c_2, c_1) = 1 = \gcd(\beta_1, \beta_1)$ *now equation 2.15 become*

$$r_2^2 = \beta_2 c_2^3 K_1 \tag{2.16}$$

from equation 2.16 $r_2^2 \mid c_2^3$ *then there exist a positive integer* e *such that* $r_2^2 e = c_2^3$ *then by using theorem 2.2*

$$c_2 = \alpha_1 \alpha_3 f^2, \quad r_2 = \alpha_3 f^3, \quad e = \alpha_1^3 \alpha_3$$

substituting the values in equation 2.16, we get $1 = \beta_2 \alpha_1^3 \alpha_3 K_1 \Longrightarrow \beta_2 = \alpha_1 = 1 = \alpha_3 = K_1$ *replacing the values we get the required result.*

Example 2.8. *Consider the 6-variable diophantine system,* $x_1 x_2 x_3 = z_1^2$ *and* $x_3 x_4 = z_2^2$*. Then all the positive solutions of this system of equations can be describe by* 11 *parametric formulas*
$x_1 = s_1^2 a d r_1 h, \quad x_2 = s_2^2 b d r_2 h, \quad x_3 = ab(c_1 r_1 r_2 dc)^2, \quad x_4 = ab(cS_1)^2,$
$z_1 = abhs_1 s_2 c c_1 r_1 r_2 d^2, \quad z_2 = abS_1 r_1 r_2 d c_1 c^2.$

Solution:-

Step 1: We apply theorem 2.4 to get the following solution for equation 1 *of the system*
$\left\{x_1 = s_1^2 k_1 h, \quad x_2 = s_2^2 l_1 h, \quad x_3 = \gamma^2 k_1 l_1, \quad z_1 = s_1 s_2 \gamma k_1 l_1 h\right\}$
and for equation 2 *of the system we get*
$\left\{x_4 = g S_1^2, \quad x_3 = S_2^2 g, \quad z_2 = g S_2 S_1\right\}$
Step 2: Since x_3 *is the repeated variable in the equations of the system, so*

$$x_3 = S_2^2 g = \gamma^2 k_1 l_1 \tag{2.17}$$

Now we will find that unique d *for* x_3*, and suppose* $a = \gcd(g, k_1)$ *such that*

$$\left\{g = a\alpha; k_1 = a\beta, \gcd(\alpha, \beta) = 1\right\} \tag{2.18}$$

then equation 2.17 becomes $S_2^2 \alpha = \gamma^2 \beta l_1 \Rightarrow \alpha \mid \gamma^2 l_1$ *so there exist a positive integer* e *such that* $\alpha e = \gamma^2 l_1$*. Now using theorem 1.3, we get* $\alpha_1 = bc^2, \quad e = b_1 c_1^2, \quad \gamma = cc_1, \quad l_1 = bb_1$ *such that* $\gcd(b, b_1) = 1 = \gcd(c, c_1)$*. Thus equation 2.17 becomes*

$S_2^2 = c_1^2\beta b_1$ clearly $c_1^2 \mid S_2^2$ and by theorem 1.2, we have $c_1 \mid S_2 \Rightarrow_2 = c_1 f$. Thus we get $f^2 = \beta b_1$ and by using theorem 2.4 we have $f = r_1 r_2 d$, $\beta = dr_1$, $b_1 = dr_2$ substituting the values we get the required result.

References

[1] W.sierpinski, Elementary Theory of Numbers, Warsaw,Poland, 1964

[2] Kennerth H. Rosen, Elementary Theory of Numbers and its applications, 5^{th} edition

[3] A.Schinzel, On the equation, $x_1 x_2 \cdots x_n = t^k$, Bill. Acad. Polon. Sci. Cl.lll, 3(1955), p.p. 17-19.

[4] M.Ward, A type of mupltication diophanitine systems, Amer.J.Math., 55(1933), pp. 67-76.

[5] Konsatantine Zelator, The diophanitine equation $xy = z^n$ for $n = 2, 3, 4, 5, 6$; The Diophanitine Equation $xyz = w^2$, and the Diophanitine systems $\{xy = v^2 \;\; yz = w^2\}$

Department of Mathematics,, National University of Computer and Emerging Sciences, B-Block, Faisal Town, Lahore, Pakistan.

E-mail address: zahid.raza@nu.edu.pk, hafsa.masood.malik@gmail.com